



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

from both the Sun and the Earth, and it is following the ephemeris very closely.

STURLA EINARSON,
A. ESTELLE GLANCY,
ALICE JOY.
BERKELEY ASTRONOMICAL DEPARTMENT,
May 19, 1907.

NOTE ON COMET *c* 1907 (GIACOBINI).

A preliminary orbit of Comet *c* 1907 has just been completed as this number of the *Publications* is going to press. The three observations used were made June 1st, 3d, and 4th, by GIACOBINI at Nice, HAMMOND at Washington, and AITKEN at Mt. Hamilton, respectively. The elements and an ephemeris are given in *Lick Observatory Bulletin*, No. 116. The observations are represented by a parabola. The inclination of the orbit plane to the ecliptic is 16° ; the longitude of the ascending node is 161° . It was nearest to the Sun on May 27th, at which time it was 117,000,000 miles distant. The comet is faint and is receding from both the Earth and the Sun. At present it is east of the sickle in *Leo*. It is traveling toward β *Leonis*, and on June 21st will be about 5° due north of it.

STURLA EINARSON,
ESTELLE GLANCY.
BERKELEY ASTRONOMICAL DEPARTMENT,
June 5, 1907.

PLANS FOR OBSERVING THE TOTAL ECLIPSE OF JANUARY, 1908.

Arrangements have been completed to dispatch an expedition from the Lick Observatory to observe the total solar eclipse of January 3, 1908. The Moon's shadow will cross the central Pacific Ocean from west to east, and pass over only two known islands. One of these is Flint Island, in longitude $151^\circ 48'$ W. and latitude $11^\circ 26'$ S.,—that is, about 390 miles northwest of the island of Tahiti.

The eclipse will occur at 11:18, local mean time, with the Sun at zenith-distance 15° . The duration given by the American Ephemeris will be $4^m 6^s$.

The generosity of Mr. WILLIAM H. CROCKER has made possible the sending of the expedition.

The expedition will sail from San Francisco on November 22d for Tahiti. The problem of transportation from Tahiti to Flint Island and return was a difficult one, but the interest of the Navy Department of the United States Government was enlisted, and the U. S. gunboat "Annapolis" has been detailed to meet the expedition at Tahiti, transport it to Flint Island,